Construct PERT network and compute:

- (i) T_E and T_L for each event
- (ii) Float for each activity
- (iii) Critical path and its duration
- (b) Solve the following LPP:

Minimum z =
$$x_1 - 3x_2 + 2X_3$$
,
Subject to, $3x_1 - X_2 + 3X_3 \le 7$
 $-2x_1 + 4x_2 \le 12$
 $-4X_1 + 3X_2 + 8X_3 \le 10$

And

$$x_1, x_2, x_3 \ge 0$$

- (c) Describe the fundamental components of a queuing process and give suitable examples.
- 7. Attempt any two questions:

12.5

(a) Find the optimal solution of the following transportation problem in which cell entries represent unit costs.

		То				
		W 1	W 2	W3	Available	
From	F1	4	14	8	10	
	F2	6	6	2	16	
	F3	10	8	14	14	
v ·	F4	2	12	4	28	
Required		14	18	36	68	

- (b) Explain the Hungarian method to solve an assignment problem.
- (c) Explain the concepts of degeneracy in simplex method.

Following Paper ID and Roll N	lo. t	o be	fill	ed i	n y	our	Ans	wer I	Book)
PAPER ID: 7113 Roll No.		I	Ι	ŀ	I	1			I
	1.B	Α.	• :					:	
(SEM. II) THEORY	EX.	\M	N.	ATIO	ON	20	10-1	1	
OPERATIO	NS	RE	SE	AR	CF	I			

Time: 3 Hours

Total Marks: 100

SECTION—A

(Attempt all questions) $(1\times10=10)$

- 1. Fill in the blanks:
 - (a) Linear programming deals with the _____. Of a function of variables known as objective function.
 - (b) The expression $a_1x_1+a_2x_2+a_3x_3+....+a_nx_n$ is ____.
 - (c) Simplex method was developed by _____.
 - (d) A set of variables is called a solution of L.P.P. if it satisfies
 - (e) Surplus variables are always accompanies with

State TRUE/FALSE:

(f) Pure strategy is the decision rule to always select a particular course of action.

 (g) Utilization factor is the average time for which a customer has to wait in the queue to get service. (h) One of the objective of the network analysis is to minimize the idle resources and investment in inventory. (i) Static models are one time decision models. (j) Vogel's approximation method is also known as regret method. (1×10=10) (i) One requirement in optimality test of transportation problem is that the number of allocations should equal to: (a) (m n) (b) (m+n+1) (c) (m+n-1) (d) None of above (ii) It is a basic feasible solution in which all the m basic variables are positive (≥ 0) and the remaining n variables are zero each: (a) Degenerate basic feasible solution (b) Non-degenerate basic feasible solution (c) Solution (d) Basic solution (iii) If the primal contains n variables and m constraints the dual will contain. (a) n variable (b) m variable (c) n+1 variable (d) m+1 variable 								
 (h) One of the objective of the network analysis is to minimize the idle resources and investment in inventory. (i) Static models are one time decision models. (j) Vogel's approximation method is also known as regret method. (1×10=10) (i) One requirement in optimality test of transportation problem is that the number of allocations should equal to: (a) (m n) (b) (m+n+1) (c) (m+n-1) (d) None of above (ii) It is a basic feasible solution in which all the m basic variables are positive (≥ 0) and the remaining n variables are zero each: (a) Degenerate basic feasible solution (b) Non-degenerate basic feasible solution (c) Solution (d) Basic solution (iii) If the primal contains n variables and m constraints the dual will contain. (a) n variable (b) m variable 	(g)	and the control of th						
 the idle resources and investment in inventory. (i) Static models are one time decision models. (j) Vogel's approximation method is also known as regret method. (1×10=10) (i) One requirement in optimality test of transportation problem is that the number of allocations should equal to: (a) (m n) (b) (m+n+1) (c) (m+n-1) (d) None of above (ii) It is a basic feasible solution in which all the m basic variables are positive (≥ 0) and the remaining n variables are zero each: (a) Degenerate basic feasible solution (b) Non-degenerate basic feasible solution (c) Solution (d) Basic solution (iii) If the primal contains n variables and m constraints the dual will contain. (a) n variable (b) m variable 	a .\							
 (i) Static models are one time decision models. (j) Vogel's approximation method is also known as regret method. (1×10=10) (i) One requirement in optimality test of transportation problem is that the number of allocations should equal to: (a) (m n) (b) (m+n+1) (c) (m+n-1) (d) None of above (ii) It is a basic feasible solution in which all the m basic variables are positive (≥ 0) and the remaining n variables are zero each: (a) Degenerate basic feasible solution (b) Non-degenerate basic feasible solution (c) Solution (d) Basic solution (iii) If the primal contains n variables and m constraints the dual will contain. (a) n variable (b) m variable 	(n)							
 (i) Vogel's approximation method is also known as regret method. (1×10=10) (i) One requirement in optimality test of transportation problem is that the number of allocations should equal to: (a) (m n) (b) (m+n+1) (c) (m+n-1) (d) None of above (ii) It is a basic feasible solution in which all the m basic variables are positive (≥ 0) and the remaining n variables are zero each: (a) Degenerate basic feasible solution (b) Non-degenerate basic feasible solution (c) Solution (d) Basic solution (iii) If the primal contains n variables and m constraints the dual will contain. (a) n variable (b) m variable 	•							
 (i) One requirement in optimality test of transportation problem is that the number of allocations should equal to: (a) (m n) (b) (m+n+1) (c) (m+n-1) (d) None of above (ii) It is a basic feasible solution in which all the m basic variables are positive (≥ 0) and the remaining n variables are zero each: (a) Degenerate basic feasible solution (b) Non-degenerate basic feasible solution (c) Solution (d) Basic solution (iii) If the primal contains n variables and m constraints the dual will contain. (a) n variable (b) m variable 	(1)							
 (i) One requirement in optimality test of transportation problem is that the number of allocations should equal to: (a) (m n) (b) (m+n+1) (c) (m+n-1) (d) None of above (ii) It is a basic feasible solution in which all the m basic variables are positive (≥ 0) and the remaining n variables are zero each: (a) Degenerate basic feasible solution (b) Non-degenerate basic feasible solution (c) Solution (d) Basic solution (iii) If the primal contains n variables and m constraints the dual will contain. (a) n variable (b) m variable 	(j)							
 (i) One requirement in optimality test of transportation problem is that the number of allocations should equal to: (a) (m n) (b) (m+n+1) (c) (m+n-1) (d) None of above (ii) It is a basic feasible solution in which all the m basic variables are positive (≥ 0) and the remaining n variables are zero each: (a) Degenerate basic feasible solution (b) Non-degenerate basic feasible solution (c) Solution (d) Basic solution (iii) If the primal contains n variables and m constraints the dual will contain. (a) n variable (b) m variable 								
 is that the number of allocations should equal to: (a) (m n) (b) (m+n+1) (c) (m+n-1) (d) None of above (ii) It is a basic feasible solution in which all the m basic variables are positive (≥ 0) and the remaining n variables are zero each: (a) Degenerate basic feasible solution (b) Non-degenerate basic feasible solution (c) Solution (d) Basic solution (iii) If the primal contains n variables and m constraints the dual will contain. (a) n variable (b) m variable 	•	(1×10=10)						
 (c) (m+n-1) (d) None of above (ii) It is a basic feasible solution in which all the m basic variables are positive (≥ 0) and the remaining n variables are zero each: (a) Degenerate basic feasible solution (b) Non-degenerate basic feasible solution (c) Solution (d) Basic solution (iii) If the primal contains n variables and m constraints the dual will contain. (a) n variable (b) m variable 	(i)							
 (ii) It is a basic feasible solution in which all the m basic variables are positive (≥ 0) and the remaining n variables are zero each: (a) Degenerate basic feasible solution (b) Non-degenerate basic feasible solution (c) Solution (d) Basic solution (iii) If the primal contains n variables and m constraints the dual will contain. (a) n variable (b) m variable 		(a) $(m n)$ (b) $(m+n+1)$						
variables are positive (≥ 0) and the remaining n variables are zero each: (a) Degenerate basic feasible solution (b) Non-degenerate basic feasible solution (c) Solution (d) Basic solution (iii) If the primal contains n variables and m constraints the dual will contain. (a) n variable (b) m variable		(c) $(m+n-1)$ (d) None of above						
 (a) Degenerate basic feasible solution (b) Non-degenerate basic feasible solution (c) Solution (d) Basic solution (iii) If the primal contains n variables and m constraints the dual will contain. (a) n variable (b) m variable 	(ii)	variables are positive (≥ 0) and the remaining n variables						
 (b) Non-degenerate basic feasible solution (c) Solution (d) Basic solution (iii) If the primal contains n variables and m constraints the dual will contain. (a) n variable (b) m variable 		are zero each:						
 (c) Solution (d) Basic solution (iii) If the primal contains n variables and m constraints the dual will contain. (a) n variable (b) m variable 		(a) Degenerate basic feasible solution						
 (d) Basic solution (iii) If the primal contains n variables and m constraints the dual will contain. (a) n variable (b) m variable 		(b) Non-degenerate basic feasible solution						
(iii) If the primal contains n variables and m constraints the dual will contain.(a) n variable(b) m variable		(c) Solution						
dual will contain. (a) n variable (b) m variable		(d) Basic solution						
(-)	(iii)							
(c) $n+1$ variable (d) $m+1$ variable		(a) n variable (b) m variable						
		(c) n+1 variable (d) m+1 variable						

(1 <u>v)</u>	func	ction of the primal a		in the of the	
	dual		•		
		Objective function			
	(b)	Non negativity restri	ctions		
	(c)	Constraints			
	(d)	All of above			
(v)		ording to this rule job decreasing due dates		sequenced in the order of	
	(a)	EDD	(b)	STR	
	(c)	SPT	(d)	FCFS	
(vi)		maximum time taken wn as:	by an	activity for completion	i
	(a)	Expected time	(b)	Optimistic time	
	(c)	Pessimistic time	(d)	None of these	
(vii)	All	the activities which a	e on th	e critical path are called	Ì
	(a)	Critical activities	(b)	Slack	
	(c)	Float	(d)	None of the above	
(viii)	In d	ecision tree analysis	lecisio	n node is represented by	,
	(a)	Circle	(b)	Triangle	
	(c)	Straight line	(d)	None of the above	
(ix)	Late	est start time can be c	alculat	ed by using:	
	(a)	Backward pass	(b)	Forward pass	
* -	(c)	Both (a) and (b)	(d)	None of the above	
(x)	The	dual of the dual is cal	lled:	•	
	(a)	Simplex	(b)	Primal	
	(c)	Second dual	(d)	None of the above	

2.

3. (i) The following table shows all the necessary information on the available supply from each warehouse, the requirement of each market and the unit transportation cost in rupees from each warehouse to each market.

Markets

		I	II	III	IV	Supply
	Α	6	3	5	4	22
	В	5	9	2	7	15
Ware house	C	5	7	8	6	8
Requirement		7	12	17	9	45

The shipping clerk has worked out the following schedule from experience: 12 units from A to II, 1 unit from A to III, 9 units from A to IV, 15 units from B to III, 7 units from C to I and 1 unit from C to III.

Find:

- (a) check if the clerk has made the optimal schedule. 5
- (b) Find the optimal schedule and minimum and minimum total shipping cost.5
- (c) If the supply from warehouse B reduces to 12 units and simultaneously the requirement of market III reduces to 14 units, find the optimal transportation schedule.
- (ii) An airline that operates 7 days a week has the time table shown below. Crew must have minimum layover of 5 hours between flights. Obtain the pairing of flights that minimizes layover time away from home assuming that the crew can

be based at either of the two cities. The crew will be based at the city that results in smaller layover.

	Delhi-	Jaipur	Jaipur-Delhi			
Fligh	nt no. Depart	Arrive	Flight no.	Depart	Arrive	
1	7 AM	8 AM	101	8 AM	9.15 AM	
2	8 AM	9 AM	102	8.30 AM	9.45 AM	
3	1.30 AM	2.30 AM	103	12 NOON	1.15 PM	
4.	6.30 PM	7.30 PM	104	5.30 PM	6.45 PM	
					15	

OR

- 3. (a) Customers arrive at the first class ticket counter of a theater at the rate of 12 per hour. There is one clerk serving the customers at the rate of 30 per hour.
 - (i) What is the probability that there is no customer in the counter?
 - (ii) What is the probability that there are more than two customers in the counter?
 - (iii) What is the probability that there is no customer waiting to be served?
 - (iv) What is the probability that a customer is being served and nobody is waiting?
 - (b) The purchase price of a machine is Rs.52,000. The installation charges amount to Rs. 14,400 and it's scrap value is only Rs. 6,400. The maintenance cost in various years is given below:

Year :1 2 3 4 5 6 7 8

Maintenance cost: 1,000 3,000 4,000 6,000 8,400 11,600 16,000 19,200

After how many year should the machine be replaced?

Assume that the machine replacement can be done only at the year ends.

3. 4. Attempt any two questions:

- 12.5
- (a) Convert the following problem into its dual:

Minimum
$$z = 2x_1 + 2x_2 + 4x_3$$
,

Subject to,
$$2x_1 + 3x_2 + 5x_3 \ge 2$$

$$3x_1 + x_2 + 7x_3 \le 3$$

$$x_1 + 4x_2 + 6x_3 \le 5$$

And $x_1, x_2, x_3 \ge 0$

(b) An investor is given the following investment alternatives and percentage rates of return.

	Low	Medium	High
Regular shares	7%	10%	15%
Risky shares	-10%	12%	25%
Property	-12%	18%	30%

Over the past 300 days, 150 days have been medium market conditions and 60 days have had high market increases.

On the basis of these data, sate the optimum investment strategy for the investment.

- (c) Give Johnson's procedure for determining an optimal sequence for processing in items on two machines. Give justification of the rule used in procedure.
- 5. Attempt any two questions:

- 12.5
- (a) Give a brief account of the methods used in model formulations.

(b) Solve the following LPP graphical method.

Maximum
$$z = x_1 + x_2$$

Subject to, $x_1 - x_2 \ge 0$

$$-3x_1 + x_2 \ge 3$$

And
$$x_1, x_2, \geq 0$$
.

(c) Solve the following game:

	. •		Player 1	В			
		I	II	Ш	IV		
	I	2	2	3	-1		
Player A	II	4	3	2	6		

6. Attempt any two questions:

12.5

(a) A project has the following times schedule:

Activity	Time in week
1-2	4
1-3	1 '
2-4	1
3-4	1
3-5	6
4-9	5
5-6	4
5-7	
6-8	1
7-8	2
8-9	1
8-10	8
9-10	7